Apply When Weeds Are Actively Growing

Sta-Green®

Phosphorus-Free

WEED & FEED

28-0-4 HERBICIDA Y FERTILIZANTE

KILLS WEEDS 2x FASTER*

FPO Spanish Kills Weeds 2 x Faster Compared to Our Previous Formula

- **Controls Dandelion, White Clover, Plantain, Chickweed, Spurge** & many other listed weeds
  Controla el diente de león, el trébol blanco, el llantén, la pamplina, la lechetrezna y muchas tras malezas enumeradas
- **Encourages Quick Greening & Provides Extended Nitrogen Feeding**
  Promueve un reverdecimiento rápido y proporciona nutrientes de nitrógeno por periodos prolongados

*Compared to our previous formula

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p acid	0.167%
Dicamba acid	0.071%
OTHER INGREDIENTS:	98.654%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
14.70 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%
3.34 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%
1.42 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%
Isomer Specific by AOAC Methods

CAS Registry Numbers:
2,4-D CAS# 1928-43-4
Mecoprop-p CAS# 16484-77-8
Dicamba CAS# 1918-00-9

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements and First Aid.

MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION Consulte el panel posterior para obtener información adicional sobre las precauciones y los primeros auxilios.

MADE IN USA

MONEY BACK Guarantee MONEY BACK

KILLS OVER 175 WEED TYPES AS LISTED

Net Wt 42 lb (19.0 kg)

Sta-Green®

Phosphorus-Free
WEED & FEED
28-0-4
KILLS WEEDS 2x FASTER*

NET WT 42 lb (19.0 kg) • COVERS 15,000 sq ft • APPROXIMATELY 1/3 ACRE

FPO UPC
0 13013 60091 4

COVERS 15,000 sq ft
APPROXIMATELY 1/3 ACRE

Sta-Green
Phosphorus-Free WEED & FEED 28-0-4
KILLS WEEDS 2x FASTER*
COVERS 15,000 sq ft APPROXIMATELY 1/3 ACRE
FPO UPC 0 13013 60091 4

Sta-Green®

THE STA-GREEN® PROMISE

YOUR SATISFACTION GUARANTEED OR YOUR MONEY BACK!

If you use Sta-Green Phosphorus-Free Weed & Feed 28-0-4 with slow-release nitrogen as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid.

STOP READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
Sta-Green Phosphorus-Free Weed & Feed 28-0-4 is a combination of three proven weed killers – 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS:

Do not use on 'Floratam' St. Augustinegrass, dichondra, carpetgrass, creeping bentgrass, or turfgrass where desirable clovers are present.
Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed the specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass (not 'Floratam'), centipedegrass, hybrid bermudagrass and colonial bentgrass.
Apply 2.8 lb of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.
Typically, a lawn utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.78 lb of nitrogen per 1,000 sq ft. Contact your local Extension Service for information specific to your geographic region.

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.
Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
Possible re-treatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Weeds Controlled

Amaranth (purple, slender); Aster (heath, white heath, purple, white prairie); Artichoke; Austrian fieldcress; Bedstraw (Catchweed, smooth, Northern, pineywoods); Beggarticks, hairy; Betony, Florida; Bindweed (Field, Hedge); Bitter Wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black Medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup (bulbous, creeping); Buttonweed (common (poorjoe), virginia); Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed (common, mousear, sticky); Chicory; Cinquefoil; Clover (crimson, alyce, hop, rabbit foot, red, strawberry, yellow sweet, white sweet, white); Cockle; Cocklebur (oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly.
NEVER APPLY BY HAND.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers 15,000 sq ft.

***SINGLE PASS* – For rotary spreaders:** Operate at intervals corresponding to the swath width. **For drop spreaders:** Overlap the wheel tracks to ensure complete coverage. For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owners' Manual.



_____ Length
x_____ Width
=_____ Total sq ft
-_____ Non-lawn areas
=_____ Total sq ft coverage to be treated

SPREADER SETTINGS

BROADCAST/ROTARY SPREADERS

Scotts® Broadcast/Rotary Spreaders Scotts Basic™, Deluxe™ EdgeGuard® Lawn Pro™, SpeedyGreen® or Standard™	5¼
Earthway EV-N-SPRED® 2050P	10

DROP SPREADERS

Scotts Drop Spreaders Scotts AccuGreen®	6

Calibrate your spreader with each application.
CALIBRATION INSTRUCTIONS: For 2.8 pounds of product per 1,000 sq ft of turfgrass: Check your spreader on 357 sq ft and adjust to apply at the rate of 1 pound of product per 357 sq ft.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
CONTAINER DISPOSAL:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-341-9620 for emergency medical treatment information.

Sta-Green Turf Note

Control Those Pesky Weeds

Many broadleaf weeds (dandelion, chickweed, henbit) germinate in the fall but are not visible in your lawn until spring. Applying Sta-Green Phosphorus-Free Weed & Feed 28-0-4 in the fall can provide effective results, because smaller weeds are easier to control. If needed, you can make a follow-up application in the spring.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water.
Do not contaminate water when disposing of equipment washwater.
When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.
Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.regulatory-info-sb.com.**

GUARANTEED ANALYSIS 28-0-4

Total Nitrogen (N)..... 28.0%
0.8% Ammoniacal Nitrogen
27.2% Urea Nitrogen*
Soluble Potash (K_2O)..... 4.0%

Derived from: Polymer-coated Urea, Urea and Potassium Chloride.
*Contains 5.7% slowly available Urea Nitrogen from coated Urea.

F1074

Manufactured for Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 2217-884-8845
EPA Est. Nos. 9198-OH-1(M), 9198-OH-2 (B), 9198-AL-1(A), 49991-TX-1 (AZ), 49991-TX-2 (BC), 45065-MN-1 (BD), 59144-GA-1(BE), 6665-NY-1(BF), 35512-FL-2(BG), 8378-IN-1(AS), 43854-MO-1(BH), 49710-TX-1(AT), 49710-TX-3(AU), 3234-UT-1(AV), 3234-AZ-1 (AW), 74764-CA-1(BK), 74764-FL-1(BL), 9404-FL-1(AX), 35497-OR-1(AY), 8660-AL-1(BN), 8660-OH-1(BO)
Circled letter is first letter of lot number. 11-10749 ©2007 UIC

www.sta-green.com
LOOK!
for other quality Sta-Green lawn & garden products. For more information, visit us on our website at www.sta-green.com or call our Consumer Service Hotline at 1.877.341.9620.
¡BUSQUE!
otros productos para césped y jardín de calidad de Sta-Green. Para obtener más información, visite nuestro sitio web www.sta-green.com o llame a nuestra línea de atención al cliente al 1.877.341.9620.
1-877-341-9620

LA PROMESA DE STA-GREEN®

¡SATISFACCION GARANTIZADA O LA DEVOLUCION DE SU DINERO!

Si aplica el herbicida y fertilizante 28-0-4 con nitrógeno de liberación continua de Sta-Green de acuerdo con las instrucciones y no queda completamente satisfecho, envíe el comprobante de compra de este paquete junto con el recibo original de la caja registradora a la dirección que aparece más abajo. Le reembolsaremos el costo de su compra.

ALTO LEA TODA LA ETIQUETA PRIMERO.
CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.
No permita que las personas (aparte de la persona que aplica el producto) ni las mascotas entren en el área de tratamiento durante la aplicación. No entre en las áreas de tratamiento hasta que el polvo se haya asentado. No se debe usar en césped que se cultive para la venta u otro uso comercial como panes de césped, ni para la producción de semillas comerciales, ni con fines de investigación.
El herbicida y fertilizante 28-0-4 de Sta-Green es una combinación de tres herbicidas probados: 2,4-D, mecoprop-p y dicamba. Este herbicida controla una amplia gama de malezas de césped, que incluyen diente de león, pamplina, polígono, llantén, ortiga mansa, lechetrezna y muchas otras que se encuentran habitualmente en céspedes residenciales. Además, un fertilizante 28-0-4 proporciona nutrientes para plantas.

PRECAUCIONES DE USO:

No lo use en las variedades 'Floratam' del césped San Agustín, dichondra, césped alfombra, "creeping bentgrass," o césped "turfgrass" cuando haya tréboles deseables.
No aplique este producto a céspedes recién cultivados hasta que estén bien establecidos o hasta después del segundo o tercer corte.
No use este producto en flores, vegetales, tapices vegetales, jardines ni plantaciones de jardín, incluidos arbustos y árboles. No aplique una dosis mayor a la especificada en ninguna zona y no aplique este producto dentro de la franja de goteo, ni sobre raíces superficiales de árboles, arbustos y otras plantas ornamentales.
Las áreas se pueden volver a sembrar o sobresembrar tres (3) semanas después de aplicar este producto.

CUANTO APLICAR:

Para usar en: Pasto azul de Kentucky, festuca spp., raigrás perenne, césped bermuda común, césped zoysia, césped San Agustín (no 'Floratam'), césped ciempiés, césped Bermuda híbrido y "colonial bentgrass."
Aplique 2.8 libras del producto cada 1,000 pies cuadrados de césped. Se puede realizar una segunda aplicación a los 30 días aproximadamente. La cantidad máxima de aplicaciones al voleo por lugar de tratamiento es de 2 por año.
En general, un césped utiliza entre 1 a 4 lb de nitrógeno cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica de acuerdo con las instrucciones, este producto proporciona 0.78 lb de nitrógeno cada 1,000 pies cuadrados. Comuníquese con la oficina local del servicio de extensión agrícola para obtener información específica para la región donde vive.

CUANDO APLICAR:

Aplique cuando las malezas de hoja ancha estén en la etapa de crecimiento activo en primavera o principios de otoño.
Aplique cuando no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, vegetales y arbustos ornamentales.
Puede ser necesario volver a aplicar el tratamiento si el césped está muy invadido por malezas, si predominan condiciones de poca humedad o si las malezas tienen poco crecimiento.

INSTRUCCIONES DE APLICACION:

1. Corte el césped hasta la altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped por lo menos 1 a 2 días antes de la aplicación para mantener la humedad hasta el siguiente riego (consulte el paso 5).
3. Aplique cuando las malezas de hoja ancha sean jóvenes y se encuentren en la etapa de crecimiento activo, preferentemente en la mañana, cuando hay rocío sobre el pasto. En el momento de la aplicación, la humedad del rocío, la lluvia o el riego/irrigación sobre las hojas de maleza, ayuda al control. Si el pasto no está húmedo cuando se hace la aplicación, rocíe ligeramente con agua antes de aplicar el producto.
4. Evite aplicar cuando las temperaturas estén constantemente por encima de los 90ºF.
5. No moje las hojas de las malezas durante 1 ó 2 días después de la aplicación. En este momento se debe regar bien.

Malezas controladas

Amaranth (purple, slender); Aster (heath, white heath, purple, white prairie); Artichoke; Austrian fieldcress; Bedstraw (Catchweed, smooth, Northern, pineywoods); Beggarticks, hairy; Betony, Florida; Bindweed (Field, Hedge); Bitter Wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black Medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup (bulbous, creeping); Buttonweed (common (poorjoe), virginia); Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed (common, mousear, sticky); Chicory; Cinquefoil; Clover (crimson, alyce, hop, rabbit foot, red, strawberry, yellow sweet, white sweet, white); Cockle; Cocklebur (oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

ESPARCIDORES Y CALIBRACION

Aplique siempre con un esparcidor giratorio o por gravedad para distribuir los gránulos uniformemente.
NUNCA APLICAR CON LA MANO.

Cobertura: Para determinar la cantidad que necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale a la superficie total en pies cuadrados. Reste todas las áreas que no sean de césped (casa, jardín, etc.). Una bolsa alcanza para cubrir 15,000 pies cuadrados.

***PASO UNICO* – Para esparcidores giratorios:** Haga funcionar a intervalos correspondientes al ancho de la banda de aspersión. **Para esparcidores por gravedad:** Superponga los carriles para asegurar una cobertura completa. Para obtener mejores resultados, aplique este producto uniformemente y evite líneas, espacios vacíos o superposiciones. Camine a ritmo regular para asegurar una cobertura uniforme. Cierre las aberturas del esparcidor cuando gire para cambiar de dirección o al detenerse.

IMPORTANTE: Es importante seguir las instrucciones según se indican en este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede ocasionar daños a su césped. Los ajustes del esparcidor son aproximados. Puede haber diferencias como consecuencia del estado del esparcidor, la velocidad a la cual se aplica y la irregularidad del área tratada. Asegúrese de examinar el esparcidor antes de comenzar y ajustarlo, si es necesario. Consulte el Manual del propietario.

_____ Largo
x_____ Ancho
=_____ Pies cuadrados totales
-_____ Áreas que no son de césped
=_____ Cobertura total en pies cuadrados a tratar

AJUSTES DEL ESPARCIDOR

ESPARCIDORES AL VOLEO/GIRATORIOS

Scotts® Broadcast/Rotary Spreaders Scotts Basic™, Deluxe™ EdgeGuard® Lawn Pro™, SpeedyGreen® or Standard™	5¼
Earthway EV-N-SPRED® 2050P	10

ESPARCIDORES POR GRAVEDAD

Scotts Drop Spreaders Scotts AccuGreen®	6

Calibre su esparcidor con cada aplicación.
INSTRUCCIONES PARA LA CALIBRACION: Para 2.8 libras de producto cada 1,000 pies cuadrados de césped: verifique que su esparcidor esté regulado en 357 pies cuadrados y ajuste para aplicar la dosis de 1 libra de producto para 357 pies cuadrados.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO DEL PESTICIDA: Guarde en el envase original y conserve todas las bolsas que contengan el producto sin usar en un área de almacenamiento cerrada, inaccesible a los niños y a los animales domésticos.
ELIMINACION DEL ENVASE:
Si está vacío: No vuelva a utilizar este envase. Tírelo en la basura, u ofrézcalo para reciclar, si existe esa opción.
Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca vierta el producto sin usar por drenajes interiores ni exteriores.

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos

PRECAUCION: Causa irritación moderada a los ojos. Nocivo si se ingiere. Evite el contacto con los ojos o la ropa. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines y zapatos. Lávese bien con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle o usar tabaco.

Primeros Auxilios

Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame a un centro de control de envenenamientos o a un médico inmediatamente para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • No administre nada por la boca a una persona que haya perdido el conocimiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-877-341-9620 para obtener información sobre tratamiento médico de urgencia.

Sta-Green Nota sobre el césped

Controla esas molestas malezas

Muchas malezas de hoja ancha (diente de león, pamplina, ortiga mansa) germinan en el otoño pero no se ven en su césped hasta la primavera. La aplicación del herbicida y fertilizante 28-0-4 de Sta-Green en el otoño puede proporcionar resultados eficaces porque las malezas pequeñas son más fáciles de controlar. Si es necesario, puede hacer otra aplicación complementaria en la primavera.

RIESGOS PARA EL MEDIO AMBIENTE:

Este producto es tóxico para los invertebrados acuáticos. El arrastre y el escurrimiento pueden afectar de manera adversa a los invertebrados acuáticos y las plantas que no sean el objetivo. No aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo.
Cuando limpie el equipo, no vierta las aguas de lavado en la tierra; rocíe o desagüe sobre un área grande lejos de pozos de agua y otras fuentes de agua.

Cuando se aplica según las instrucciones, este producto cumple con las normas para metales aceptadas por la Association of American Plant Food Control Officials.
Se puede obtener información con respecto al contenido y los niveles de metales de este producto en la siguiente dirección de Internet:
http://www.regulatory-info-sb.com.

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Net Wt 42 lb (19.0 kg)

PROOF OF PURCHASE
COMPROBANTE DE COMPRA

FPO UPC
0 13013 60091 4

COVERS 15,000 sq ft • APPROXIMATELY 1/3 ACRE

Sta-Green
Phosphorus-Free
WEED & FEED
28-0-4
KILLS WEEDS 2X FASTER*
COVERS 15,000 sq ft
APPROXIMATELY 1/3 ACRE
FPO UPC
0 13013 60091 4
Apply When Weeds Are Actively Growing

K FASTER*

COVERS 15,000 sq ft
APPROXIMATELY 1/3 ACRE

FPO UPC
0 13013 60091 4

28-0-4 HERBICIDA Y FERTILIZANTE

KILLS WEEDS 2x FASTER*

FPO Spanish Kills Weeds 2 x Faster Compared to Our Previous Formula

- **Controls Dandelion, White Clover, Plantain, Chickweed, Spurge** & many other listed weeds
  Controla el diente de león, el trébol blanco, el llantén, la pamplina, la lechetrezna y muchas tras malezas enumeradas
- **Encourages Quick Greening & Provides Extended Nitrogen Feeding**
  Promueve un reverdecimiento rápido y proporciona nutrientes de nitrógeno por periodos prolongados

Phosphorus-Free
WEED & FEED

Sta-Green®

*Compared to our previous formula

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p acid	0.167%
Dicamba acid	0.071%
OTHER INGREDIENTS:	98.654%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
14.70 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%
3.34 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%
1.42 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%
Isomer Specific by AOAC Methods

CAS Registry Numbers:
2,4-D CAS# 1928-43-4
Mecoprop-p CAS# 16484-77-8
Dicamba CAS# 1918-00-9

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements and First Aid.

MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION Consulte el panel posterior para obtener información adicional sobre las precauciones y los primeros auxilios.

MADE IN USA

MONEY BACK Guarantee MONEY BACK

KILLS OVER 175 WEED TYPES AS LISTED

Net Wt 42 lb (19.0 kg)

Sta-Green®

Phosphorus-Free
WEED & FEED
28-0-4

KILLS WEEDS 2x FASTER*

NET WT 42 lb (19.0 kg) • COVERS 15,000 sq ft • APPROXIMATELY 1/3 ACRE

THE STA-GREEN® PROMISE

YOUR SATISFACTION GUARANTEED OR YOUR MONEY BACK!

If you use Sta-Green Phosphorus-Free Weed & Feed 28-0-4 with slow-release nitrogen as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid.

STOP READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
Sta-Green Phosphorus-Free Weed & Feed 28-0-4 is a combination of three proven weed killers – 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS:

Do not use on 'Floratam' St. Augustinegrass, dichondra, carpetgrass, creeping bentgrass, or turfgrass where desirable clovers are present.
Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed the specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass (not 'Floratam'), centipedegrass, hybrid bermudagrass and colonial bentgrass.
Apply 2.8 lb of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.
Typically, a lawn utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.78 lb of nitrogen per 1,000 sq ft. Contact your local Extension Service for information specific to your geographic region.

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.
Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
Possible re-treatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Weeds Controlled

Amaranth (purple, slender); Aster (heath, white heath, purple, white prairie); Artichoke; Austrian fieldcress; Bedstraw (Catchweed, smooth, Northern, pineywoods); Beggarticks, hairy; Betony, Florida; Bindweed (Field, Hedge); Bitter Wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black Medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup (bulbous, creeping); Buttonweed (common (poorjoe), virginia); Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed (common, mouseear, sticky); Chicory; Cinquefoil; Clover (crimson, alyce, hop, rabbit foot, red, strawberry, yellow sweet, white sweet, white); Cockle; Cocklebur (oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly.

SPREADER SETTINGS

BROADCAST/ROTARY SPREADERS

LA PROMESA DE STA-GREEN®

¡SATISFACCION GARANTIZADA O LA DEVOLUCION DE SU DINERO!

Si aplica el herbicida y fertilizante 28-0-4 con nitrógeno de liberación continua de Sta-Green de acuerdo con las instrucciones y no queda completamente satisfecho, envíe el comprobante de compra de este paquete junto con el recibo original de la caja registradora a la dirección que aparece más abajo. Le reembolsaremos el costo de su compra.

ALTO LEA TODA LA ETIQUETA PRIMERO.
CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.
No permita que las personas (aparte de la persona que aplica el producto) ni las mascotas entren en el área de tratamiento durante la aplicación. No entre en las áreas de tratamiento hasta que el polvo se haya asentado. No se debe usar en césped que se cultive para la venta u otro uso comercial como panes de césped, ni para la producción de semillas comerciales, ni con fines de investigación.
El herbicida y fertilizante 28-0-4 de Sta-Green es una combinación de tres herbicidas probados: 2,4-D, mecoprop-p y dicamba. Este herbicida controla una amplia gama de malezas de césped, que incluyen diente de león, pamplina, polígono, llantén, ortiga mansa, lechetrezna y muchas otras que se encuentran habitualmente en céspedes residenciales. Además, un fertilizante 28-0-4 proporciona nutrientes para plantas.

PRECAUCIONES DE USO:

No lo use en las variedades 'Floratam' del césped San Agustín, dichondra, césped alfombra, "creeping bentgrass," o césped "turfgrass" cuando haya tréboles deseables.
No aplique este producto a céspedes recién cultivados hasta que estén bien establecidos o hasta después del segundo o tercer corte.
No use este producto en flores, vegetales, tapices vegetales, jardines ni plantaciones de jardín, incluidos arbustos y árboles. No aplique una dosis mayor a la especificada en ninguna zona y no aplique este producto dentro de la franja de goteo, ni sobre raíces superficiales de árboles, arbustos y otras plantas ornamentales.
Las áreas se pueden volver a sembrar o sobresembrar tres (3) semanas después de aplicar este producto.

CUANTO APLICAR:

Para usar en: Pasto azul de Kentucky, festuca spp., raigrás perenne, césped bermuda común, césped zoysia, césped San Agustín (no 'Floratam'), césped ciempiés, césped Bermuda híbrido y "colonial bentgrass."
Aplique 2.8 libras del producto cada 1,000 pies cuadrados de césped. Se puede realizar una segunda aplicación a los 30 días aproximadamente. La cantidad máxima de aplicaciones al voleo por lugar de tratamiento es de 2 por año.
En general, un césped utiliza entre 1 a 4 lb de nitrógeno cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica de acuerdo con las instrucciones, este producto proporciona 0.78 lb de nitrógeno cada 1,000 pies cuadrados. Comuníquese con la oficina local del servicio de extensión agrícola para obtener información específica para la región donde vive.

CUANDO APLICAR:

Aplique cuando las malezas de hoja ancha estén en la etapa de crecimiento activo en primavera o principios de otoño.
Aplique cuando no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, vegetales y arbustos ornamentales.
Puede ser necesario volver a aplicar el tratamiento si el césped está muy invadido por malezas, si predominan condiciones de poca humedad o si las malezas tienen poco crecimiento.

INSTRUCCIONES DE APLICACION:

1. Corte el césped hasta la altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped por lo menos 1 a 2 días antes de la aplicación para mantener la humedad hasta el siguiente riego (consulte el paso 5).
3. Aplique cuando las malezas de hoja ancha sean jóvenes y se encuentren en la etapa de crecimiento activo, preferentemente en la mañana, cuando hay rocío sobre el pasto. En el momento de la aplicación, la humedad del rocío, la lluvia o el riego/irrigación sobre las hojas de maleza, ayuda al control. Si el pasto no está húmedo cuando se hace la aplicación, rocíe ligeramente con agua antes de aplicar el producto.
4. Evite aplicar cuando las temperaturas estén constantemente por encima de los 90ºF.
5. No moje las hojas de las malezas durante 1 ó 2 días después de la aplicación. En este momento se debe regar bien.

Malezas controladas

Amaranth (purple, slender); Aster (heath, white heath, purple, white prairie); Artichoke; Austrian fieldcress; Bedstraw (Catchweed, smooth, Northern, pineywoods); Beggarticks, hairy; Betony, Florida; Bindweed (Field, Hedge); Bitter Wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black Medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup (bulbous, creeping); Buttonweed (common (poorjoe), virginia); Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed (common, mouseear, sticky); Chicory; Cinquefoil; Clover (crimson, alyce, hop, rabbit foot, red, strawberry, yellow sweet, white sweet, white); Cockle; Cocklebur (oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

ESPARCIDORES Y CALIBRACION

Aplique siempre con un esparcidor giratorio o por gravedad para distribuir los gránulos uniformemente.

AJUSTES DEL ESPARCIDOR

ESPARCIDORES AL VOLEO/GIRATORIOS

(oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly.
NEVER APPLY BY HAND.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers 15,000 sq ft.

***SINGLE PASS* – For rotary spreaders:** Operate at intervals corresponding to the swath width. **For drop spreaders:** Overlap the wheel tracks to ensure complete coverage. For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owners' Manual.



_____ Length
x_____ Width
=_____ Total sq ft
- _____ Non-lawn areas
=_____ Total sq ft coverage to be treated

SPREADER SETTINGS

BROADCAST/ROTARY SPREADERS

Scotts® Broadcast/Rotary Spreaders Scotts Basic™, Deluxe™ EdgeGuard® Lawn Pro™, SpeedyGreen® or Standard™	5¼
Earthway EV-N-SPRED® 2050P	10

DROP SPREADERS

Scotts Drop Spreaders Scotts AccuGreen®	6

Calibrate your spreader with each application.
CALIBRATION INSTRUCTIONS: For 2.8 pounds of product per 1,000 sq ft of turfgrass: Check your spreader on 357 sq ft and adjust to apply at the rate of 1 pound of product per 357 sq ft.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
CONTAINER DISPOSAL:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-341-9620 for emergency medical treatment information.	

Sta-Green Turf Note

Control Those Pesky Weeds

Many broadleaf weeds (dandelion, chickweed, henbit) germinate in the fall but are not visible in your lawn until spring. Applying Sta-Green Phosphorus-Free Weed & Feed 28-0-4 in the fall can provide effective results, because smaller weeds are easier to control. If needed, you can make a follow-up application in the spring.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.
When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.
Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.regulatory-info-sb.com.**

GUARANTEED ANALYSIS 28-0-4

Total Nitrogen (N)..... 28.0%
0.8% Ammoniacal Nitrogen
27.2% Urea Nitrogen*
Soluble Potash (K_2O)..... 4.0%

Derived from: Polymer-coated Urea, Urea and Potassium Chloride.
*Contains 5.7% slowly available Urea Nitrogen from coated Urea.

F1074

Manufactured for Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 2217-884-8845
EPA Est. Nos. 9198-OH-1(M), 9198-OH-2(B), 9198-AL-1(A), 49991-TX-1(AZ), 49991-TX-2(BC), 45065-MN-1(BD), 59144-GA-1(BE), 6665-NY-1(BF), 35512-FL-2(BG), 8378-IN-1(AS), 43854-MO-1(BH), 49710-TX-1(AT), 49710-TX-3(AU), 3234-UT-1(AV), 3234-AZ-1(AW), 74764-CA-1(BK), 74764-FL-1(BL), 9404-FL-1(AX), 35497-OR-1(AY), 8660-AL-1(BN), 8660-OH-1(BO)
Circled letter is first letter of lot number. 11-10749 ©2007 UIC

www.sta-green.com
LOOK! for other quality Sta-Green lawn & garden products. For more information, visit us on our website at www.sta-green.com or call our Consumer Service Hotline at 1.877.341.9620.
¡BUSQUE! otros productos para césped y jardín de calidad de Sta-Green. Para obtener más información, visite nuestro sitio web www.sta-green.com o llame a nuestra línea de atención al cliente al 1.877.341.9620.
1-877-341-9620

Riesgos para los seres humanos y los animales domésticos

PRECAUCION: Causa irritación moderada a los ojos. Nocivo si se ingiere. Evite el contacto con los ojos o la ropa. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines y zapatos. Lávese bien con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle o usar tabaco.

Primeros Auxilios	
Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame a un centro de control de envenenamientos o a un médico inmediatamente para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • No administre nada por la boca a una persona que haya perdido el conocimiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-877-341-9620 para obtener información sobre tratamiento médico de urgencia.	

Sta-Green Nota sobre el césped

Controla esas molestas malezas

Muchas malezas de hoja ancha (diente de león, pamplina, ortiga mansa) germinan en el otoño pero no se ven en su césped hasta la primavera. La aplicación del herbicida y fertilizante 28-0-4 de Sta-Green en el otoño puede proporcionar resultados eficaces porque las malezas pequeñas son más fáciles de controlar. Si es necesario, puede hacer otra aplicación complementaria en la primavera.

RIESGOS PARA EL MEDIO AMBIENTE:

Este producto es tóxico para los invertebrados acuáticos. El arrastre y el escurrimiento pueden afectar de manera adversa a los invertebrados acuáticos y las plantas que no sean el objetivo. No aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo.
Cuando limpie el equipo, no vierta las aguas de lavado en la tierra; rocíe o desagüe sobre un área grande lejos de pozos de agua y otras fuentes de agua.

Cuando se aplica según las instrucciones, este producto cumple con las normas para metales aceptadas por la Association of American Plant Food Control Officials.
Se puede obtener información con respecto al contenido y los niveles de metales de este producto en la siguiente dirección de Internet:
http://www.regulatory-info-sb.com.

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Net Wt 42 lb (19.0 kg)

PROOF OF PURCHASE
COMPROBANTE DE COMPRA
FPO UPC
0 13013 60091 4

(oriental, spiny); Common mullein; Corn Spurry; Creeping Beggarweed; Creeping Jenny (creeping charlie); Cupid shaving Brush; Cudweed; Daisy (english, oxeye); Dandelion (common, false); Day flower; Deadnettle; Dichondra; Dock (broadleaf, curly); Dogbane; Dogfennel; Dollarweed (Pennywort); Dove weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree (redstem, whitestem); Fleabane (daisy (annual), rough); Galinsoga (hairy, smallflower); Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed (meadow, mouseear, orange, yellow); Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-Jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow (alkali, bristly, common, dwarf, venice); Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory (bigroot, ivyleaf, mexican, red, tall, wolly); Mugwort; Mustard (ball, black, blue, hedge, indian, tansy, tumble, white, wormseed); Nettle (stinging (burning), tall, wood); Parsley-piert; Parsnip; Pearlwort; Pennycress (frenchweed); Pepperweed (perennial, virginia (Peppergrass)); Pigweed (prostate, redroot, russian, smooth, tumble); Pineapple weed; Plains coreopsis (tickseed); Plantain (blackseed, bracted, broadleaf, buckhorn, hoary, narrowleaf, slender, woolly); Poison ivy; Poison oak; Pokeweed; Prairie Sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley (brazil, florida, large flower); Ragweed (bur, common, lancleaf, western); Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepardspurse; Smartweed (ladysthumb, pale, pennsylvania); Smooth chaff-flower; Smoothdock; Sorrel (red (sheep)); Sowthistle (annual, spiny); Spanish needles; Speedwell (bi-lobed, common, corn, ivyleaf, persian, purslane, slender, snow, thymeleaf); Spiny amaranth; Spurge (spotted, prostrate); Spurweed; Star of Bethlehem; Strawberry, India mock; Tansy ragwort; Tanweed; Thistle (blessed, bull, canada, flodman (prairie), musk, platte, plumeless, russian, scotch, wavyleaf, yellow (horrible)); Trailing crownvetch; Velvetleaf; Vervain (blue, hoary, prostrate, tall); Vetch (bird, hairy, milk, narrowleaf, wild); Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckweat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel (creeping, violet, yellow (Oxalis)); Woolly croton; Wormseed; Yarrow (common, western, yellow)

ESPARCIDORES Y CALIBRACION

Aplique siempre con un esparcidor giratorio o por gravedad para distribuir los gránulos uniformemente.
NUNCA APLICAR CON LA MANO.

Cobertura: Para determinar la cantidad que necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale a la superficie total en pies cuadrados. Reste todas las áreas que no sean de césped (casa, jardín, etc.). Una bolsa alcanza para cubrir 15,000 pies cuadrados.

***PASO UNICO* – Para esparcidores giratorios:** Haga funcionar a intervalos correspondientes al ancho de la banda de aspersión. **Para esparcidores por gravedad:** Superponga los carriles para asegurar una cobertura completa. Para obtener mejores resultados, aplique este producto uniformemente y evite líneas, espacios vacíos o superposiciones. Camine a ritmo regular para asegurar una cobertura uniforme. Cierre las aberturas del esparcidor cuando gire para cambiar de dirección o al detenerse.

IMPORTANTE: Es importante seguir las instrucciones según se indican en este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede ocasionar daños a su césped. Los ajustes del esparcidor son aproximados. Puede haber diferencias como consecuencia del estado del esparcidor, la velocidad a la cual se aplica y la irregularidad del área tratada. Asegúrese de examinar el esparcidor antes de comenzar y ajustarlo, si es necesario. Consulte el Manual del propietario.



_____ Largo
x_____ Ancho
=_____ Pies cuadrados totales
- _____ Áreas que no son de césped
=_____ Cobertura total en pies cuadrados a tratar

AJUSTES DEL ESPARCIDOR

ESPARCIDORES AL VOLEO/GIRATORIOS

Scotts® Broadcast/Rotary Spreaders Scotts Basic™, Deluxe™ EdgeGuard® Lawn Pro™, SpeedyGreen® or Standard™	5¼
Earthway EV-N-SPRED® 2050P	10

ESPARCIDORES POR GRAVEDAD

Scotts Drop Spreaders Scotts AccuGreen®	6

Calibre su esparcidor con cada aplicación.
INSTRUCCIONES PARA LA CALIBRACION: Para 2.8 libras de producto cada 1,000 pies cuadrados de césped: verifique que su esparcidor esté regulado en 357 pies cuadrados y ajuste para aplicar la dosis de 1 libra de producto para 357 pies cuadrados.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO DEL PESTICIDA: Guarde en el envase original y conserve todas las bolsas que contengan el producto sin usar en un área de almacenamiento cerrada, inaccesible a los niños y a los animales domésticos.
ELIMINACION DEL ENVASE:
Si está vacío: No vuelva a utilizar este envase. Tírelo en la basura, u ofrézcalo para reciclar, si existe esa opción.
Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca vierta el producto sin usar por drenajes interiores ni exteriores.

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos

PRECAUCION: Causa irritación moderada a los ojos. Nocivo si se ingiere. Evite el contacto con los ojos o la ropa. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines y zapatos. Lávese bien con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle o usar tabaco.

Primeros Auxilios	
Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame a un centro de control de envenenamientos o a un médico inmediatamente para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • No administre nada por la boca a una persona que haya perdido el conocimiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-877-341-9620 para obtener información sobre tratamiento médico de urgencia.	

Sta-Green Nota sobre el césped

Controla esas molestas malezas

Muchas malezas de hoja ancha (diente de león, pamplina, ortiga mansa) germinan en el otoño pero no se ven en su césped hasta la primavera. La aplicación del herbicida y fertilizante 28-0-4 de Sta-Green en el otoño puede proporcionar resultados eficaces porque las malezas pequeñas son más fáciles de controlar. Si es necesario, puede hacer otra aplicación complementaria en la primavera.

COVERS 15,000 sq ft • APPROXIMATELY 1/3 ACRE